

SYNCHRONIZED RF & HIFEM: BUTTOCK MRI STUDY

COMBINED HIFEM AND SYNCHRONIZED RADIOFREQUENCY ENERGIES ACHIEVE SUPERIOR GLUTEAL MUSCLE CONTOURING THAN HIFEM ENERGY ALONE

Barry DiBernardo, MD¹; David J. Goldberg, MD, JD²; Bruce Katz, MD³;
JD McCoy, NMD⁴; Suneel Chilukuri, MD, FAAD, FACMS⁵

1. New Jersey Plastic Surgery, New Jersey, NY, USA; 2. Icahn School of Medicine, New York, NY, USA; 3. Juva Skin and Laser Center, Manhattan, NY, USA; 4. Contour Medical, Gilbert, AZ, USA; 5. Refresh Dermatology, Houston, TX, USA;

Published in Aesthetic Surgery Journal Open Forum, January 2023, DOI:10.1093/asjof/ojac087

HIGHLIGHTS

- 67 subjects (21-67 years, 16-34 kg/m², skin type I-VI) were divided into HIFEM+RF group (n=34) and Standalone HIFEM group (n=33), both receiving four 30-minute therapies
- Group HIFEM+RF showed an average **+24.7% increase**, while standalone HIFEM group showed an average 15.9% increase in muscle thickness at 3 months follow-up visit
- There was no significant reduction in subcutaneous fat thickness in the gluteal area (p-value>0.05)




Figure 1: 21-year-old woman treated with combination of RF+HIFEM, digital photographs were taken at baseline (left), after the 4th treatment (middle), at 3-month follow-up (right, a 22.8% increase in muscle mass), the dotted line shows the shape of the buttocks at baseline.